



Effect of Foliar Application of Growth Retardants on Reproductive Efficiency of Groundnut (*Arachishypogaea*L.) Genotypes

O. Venkata Narayana ^{a++*}, B. Santhosh ^{b#}, T. Raghavendra ^{ct}
and M. Raghavendra ^{at}

^a S.V. Agricultural College, Tirupati, AP, India.

^b Agricultural College, Rajamahendravaram, AP, India.

^c Regional Agricultural Research Station, Nandhyal, AP, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/arja/2024/v17i4635>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129027>

Original Research Article

Received: 24/10/2024

Accepted: 28/12/2024

Published: 30/12/2024

ABSTRACT

A field experiment entitled "Study on foliar application of Plant growth regulators for improving reproductive efficiency and yield of Groundnut (*Arachishypogaea*L.) genotypes" was carried out during Rabi, 2023-24 at wetland farm of S. V. Agricultural College, Tirupati campus of Acharya N.

⁺⁺M.Sc. Research Scholar Crop Physiology;

[#]Assistant Professor (Crop Physiology);

[†]Scientist(Crop Physiology);

[‡]Associate Professor (Bio chemistry);

*Corresponding author: E-mail: b.santhosh@angrau.ac.in;

Cite as: Narayana, O. Venkata, B. Santhosh, T. Raghavendra, and M. Raghavendra. 2024. "Effect of Foliar Application of Growth Retardants on Reproductive Efficiency of Groundnut (*Arachishypogaea*L.) Genotypes". *Asian Research Journal of Agriculture* 17 (4):1202-7. <https://doi.org/10.9734/arja/2024/v17i4635>.

G. Ranga Agricultural University, Andhra Pradesh. This experiment was laid out in randomized block design with factorial concept and replicated thrice, with four genotypes as factor-1 (G₁-Nityaharitha, G₂-Dharani, G₃-K-6, G₄-Vishista) and four plant growth regulators applied on as Factor-2 (T₁- Control + RDF of NPK, T₂-Prohexadione calcium@105g a.i/ha, T₃- Putrescine@100 ppm, T₄- Paclobutrazol@ 100 ppm applied before and at peak flowering stage of crop). The reproductive efficiency attributes viz., number of flowers, pegs and pods per plant, flower to peg ratio, flower to pod ratio and length of peg were recorded and analysed to understand the influence of plant growth regulators on groundnut. G₁ recorded highest number for pegs per plant (54.6), total pods per plant (36.8), mature pods per plant (28.5) and lowest flower to peg ratio (2.8), lowest length of peg (5.0cm). whereas G₂ showed lowest flower to pod ratio (4.3). Treatment of T₂ resulted in highest number of pegs per plant (51.8), total pods per plant (36.9), mature pods per plant (29.7), lowest flower to peg ratio (2.6), minimum length of peg (5.2 cm) lowest flower to pod ratio (3.6) and lowest number of flowers per plant (134.9) compared to the other treatments. Nityaharita and Prohexadione calcium@105g a.i/ha resulted in better reproductive efficiency in groundnut.

Keywords: Groundnut; plant growth regulators; growth retardants; Paclobutrazol; reproductive efficiency.

1. INTRODUCTION

Groundnut belongs to the family fabaceae and order fabales. It is a widely cultivated annual crop that exhibits dicotyledonous behaviour and self-pollination. Groundnut is grown in tropical, sub-tropical, and warm climate zones between the latitudes of 40° N and 40° S. Groundnut is a rich source of edible oil (47-54%), high-quality protein (22-30%), starch (6-24%), cellulose (1-2%), minerals (2-3%) and calories (5-6%). Globally, Groundnut is cultivated under 327 lakh hectares, producing 539 lakh tonnes with the productivity of 1648 kg per hectare [1]. India has 54.2 Lakh hectares area under Groundnut cultivation with a productivity of 1863 kg per hectare. In Andhra Pradesh, groundnut is cultivated in 8.23 lakh hectares with a production of 5.19 lakh tones. Groundnut seed contains 44 to 56% oil and 22 to 30% protein. Kernels are rich sources of riboflavin, thiamine, nicotinic acid and vitamin E and is rich in minerals like P, Ca, Mg and K and vitamins.

Reproductive efficiency of groundnut depends primarily on light absorption, assimilates production, production of viable flowers, pegs, flower to peg ratio, conversion of and pegs to filled pods [2]. Putrescine is involved in various physiological processes, including growth, development and stress responses and results in increased crop yields [3]. Paclobutrazole is widely proven to inhibit the biosynthesis of gibberellins, which promote stem elongation, resulting in compact plants with shorter internodes with

stouter stem, increasing root growth and causing better fruit set [4]. Prohexadione Calcium controls excessive vegetative growth, resulting in shorter internodes and a more compact plant structure. It suppresses apical dominance, leading to increased branching and lateral bud development [5].

The farmers in arid regions of Andhra Pradesh struggle for finding sufficient water supply for the crop during critical stages of crop growth and suitable chemicals for enhancing the flowering window and reproductive productivity. To address the problem, the current study is taken up with an objective to understand the effect of growth retardants on reproductive efficiency of Groundnut.

2. MATERIALS AND METHODS

A field experiment entitled "Study on foliar application of plant growth regulators for improving reproductive efficiency and yield of Groundnut (*Arachis hypogaea* L.) genotypes" was conducted on sandy clay loam soils of wetland farm of S.V. Agricultural College, Tirupati, Acharya N.G. Ranga Agricultural University. The experiment was laid out in a randomized block design with factorial concept and replicated thrice. The experiment consisted of 16 combinations with four groundnut genotypes viz., Nityaharitha, Dharani, K-6 and Vishista and four growth regulator treatments viz., control, Prohexadione Calcium @105 g a.i/ha, Putrescine @ 100ppm and Paclobutrazol @100ppm.

The weekly mean temperature during the crop growth period is 33.72°C and mean relative humidity is 60.75 %. A total rainfall of 16.0 mm received during the crop growth period in 1 rainy day as against the decennial average of 92.9 mm of rainfall received in 4.5 rainy days. The decennial mean sunshine hours for the corresponding period is 7.2 hours. The experimental plots were of 3x2 meter size and spacing of 22.5 cm x 10 cm is maintained.

Pre harvest chemical treatment sprays were prepared and applied before and during flowering. In order to prepare 105 g a.i./ha Prohexadione calcium, 105 g of chemical active ingredient is dissolved in 500 litres of water and applied per hectare. In order to prepare 100 ppm of Putrescine, 100 mg of chemical was dissolved in 1 litre of water. In order to prepare 100ppm of Paclobutrazole, 100 mg of chemical is dissolved in 1 litre of water.

Number of flowers was computed by summing up the number of flowers produced per plant during entire flowering period. The number of pegs from the five plants tagged were counted and the average was expressed as number of pegs per plant. The number of pods from the five different plants tagged were counted and the average was expressed as total number of pods per plant. The number of mature pods from the tagged plants were counted and the average was expressed as number of mature pods per plant. It is the ratio of total number of flowers produced per plant to the number of pegs from the plants tagged were counted and the average was expressed as flowers to peg ratio.

It is the ratio of total number of flowers produced per plant during entire flowering period to the number of pods from the five plants tagged which is calculated as flowers to pod ratio. Length of peg(cm) of the five randomly selected and tagged plants in each plot was measured from the base of the flower to the growing point of the peg using a scale which contains numerical values in centimeters (cm). The statistical analysis of experimental data was analyzed in factorial randomised block design and OPSTAT was the statistical package used for data analysis.

3. RESULTS AND DISCUSSION

The number of flowers per plant of groundnut was significantly influenced by the genotypes. Among the genotypes of the study, the highest (158.8) and lowest (142.3) number of flowers per plant of groundnut were recorded from G₃ and G₄ respectively. There was significant difference found among genotypes and treatments. The maximum and minimum number of flowers per plant was observed with T₁ (173.7) and T₂ (134.9) respectively. And among all of the interactions of genotypes with treated plant growth regulator treatments, the highest number flowers per plant (181.7) found in G₂, treated with T₁. This treatment could result in reduced vegetative growth and translocate those resources to flowers and thereby enhancing the maximum number of flowers and seed filling percentage. The application of PBZ to peanut plants at the stage of pod formation, seed yield was increased [6].

The maximum number of pegs per plant of groundnut was recorded with the G₁ (54.6) which indicates efficiency of the genotype to transform flowers into pegs which could led to better pegs production and yield. Whereas the lowest number recorded with G₃ (42.3). Among the PGRs studied, the maximum number of pegs per plant was observed with T₂ (51.8), and minimum number of pegs per plant recorded with T₁ (44.1). The maximum (57.1) and minimum (36.2) number of pegs per plant were recorded with interaction of G₁ when sprayed with T₂ and G₃ (36.2) with T₁ among all interactions. These results are in similarity with Hua et al. [7 and 8].

The maximum number of pods per plant recorded with the G₁ (36.8) followed by G₂ (36.6) and T₂ (36.9) and lowest number of total pods per plant recorded with G₄ (25.6) and T₁ (21.0). G₁ applied with T₂ recorded highest (40.9) and G₄ when treated with T₁ recorded minimum number of total pods (21.0). The number of mature pods per plant of groundnut was recorded maximum with the G₁ (28.5), T₂ (29.7) and G₁ treated with T₂ (32.5). This could be attributed to increased pegs production and better assimilate translocation efficiency of G₁ which ultimately leads to greater mature pods production potential.

The minimum mature pods per plant recorded with G₄ (22.6) and T₁ (20.3). These results were in accordance with Gaudin [9].

Table 1. Total number of flowers, pegs and length of peg of groundnut genotypes as influenced by PGRs

Treatments	Total Number of Flowers					Total Number of Pegs					Length of Peg(cm)				
	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean
G1	171.5	137.3	159.3	148.7	154.2	51.2	57.1	54.3	55.7	54.6	5.1	4.6	5.6	4.7	5.0
G2	181.7	138.8	161.8	149.2	157.9	49.3	56.7	52.9	54.9	53.5	5.7	5.2	6.1	5.3	5.6
G3	179.0	139.5	164.4	152.1	158.8	36.2	46.6	42.0	44.2	42.3	6.1	5.6	6.5	5.8	6.0
G4	162.5	124.0	144.3	138.3	142.3	39.5	46.8	43.8	45.3	43.8	5.9	5.4	6.5	5.7	5.9
Mean	173.7	134.9	157.5	147.1		44.1	51.8	48.2	50.0		5.7	5.2	6.2	5.4	
	G	T	GXT			G	T	GXT			G	T	GXT		
SEm	3.53	3.53	7.06			0.35	0.35	0.70			0.038	0.038	0.076		
CD(P=0.05)	10.201	10.201	NS			1.01	1.01	NS			0.109	0.109	NS		

Table 2. Total number of pods per plant and mature pods per plant, flower to peg ratio and flower to pod ratio of groundnut genotypes as influenced by PGRs

Treatments	Total number of pods per plant					Total number of mature pods per plant					Flower to Peg Ratio					Flower to Pod Ratio				
	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean
G1	30.7	40.9	37.0	38.7	36.8	22.3	32.5	28.7	30.3	28.5	3.4	2.4	2.9	2.7	2.8	5.7	3.4	4.4	3.9	4.4
G2	30.0	40.0	38.2	38.3	36.6	21.0	31.7	29.5	29.3	27.9	3.7	2.5	3.1	2.7	3.0	6.0	3.4	4.2	3.8	4.3
G3	29.0	38.0	36.3	37.0	35.1	20.0	29.0	27.4	28.3	26.2	5.0	3.0	3.9	3.4	3.8	5.9	3.5	4.3	3.9	4.4
G4	21.0	28.7	25.9	26.7	25.6	18.0	25.7	23.0	23.7	22.6	4.1	2.7	3.3	3.1	3.3	7.5	4.2	5.4	5.1	5.6
Mean	27.7	36.9	34.4	35.2		20.3	29.7	27.2	27.9		4.0	2.6	3.3	3.0		6.3	3.6	4.6	4.2	
	G	T	GXT			G	T	GXT			G	T	GXT			G	T	GXT		
SEm+	0.45	0.45	0.91			0.47	0.47	0.94			0.084	0.084	0.169			0.117	0.117	0.233		
CD (P=0.05)	1.31	1.31	NS			1.36	1.36	NS			0.243	0.243	NS			0.337	0.337	NS		

The minimum of flower to peg ratio of groundnut was recorded with the G₁ (2.8), T₂ (2.6). And highest flower to peg ratio recorded in genotype G₃ (3.8), T₁ (4.0) and G₃ treated with T₁ (5.0) because of inherent genetic characteristics of genotypes.

The highest of flower to pod ratio was recorded with the G₄ (5.6), T₁ (6.3) and G₄ treated with T₁ (7.5). And lowest flower to peg ratio recorded in genotype G₂ (4.3) and T₂ (3.6).

Minimum length of peg of groundnut was recorded with the G₁ (5.0 cm) and T₂ (5.2 cm). Maximum length of peg recorded in genotype G₃ (6.0 cm), T₃ (6.2 cm) which increases plant height by cell division, cell elongation and cell differentiation due to induction of auxins and cytokinin. By treatment with TIBA, peg length inhibited by 25.08% than the control due to reduced plant height results in pegs travel less distance to reach earth surface [10].

4. CONCLUSION

The foliar application of Prohexadione Calcium, Paclobutrazol in at specified concentrations applied to groundnut before flowering stage, had been proved better in reducing excessive vegetative growth and increased the flower and pod number of groundnut. K-6 have been reported better number of flowers and flower to pod ratio. The present study concludes that, the reproductive efficiency obtained with Prohexadione Calcium@ 105ga.i/hain Groundnut, on sandy clay loam soils of Southern Agro-climatic Zone of Andhra Pradesh during Rabi season.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

ACKNOWLEDGEMENTS

Authors are grateful to the ANGRAU and Department of crop physiology, S.V. Agricultural College, Tirupati for providing essential facilities and laboratory equipment to conduct this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAOSTAT, Food and Agricultural Organization of United Nations Database; 2021.
2. Swethasree M, Sudhakar P, Umamahesh V, Prathima T, Giridhara TK. Effect of planting density on yield and architecture suitability of groundnut (*Arachis hypogaea* L.) varieties. The Indian Journal of Agricultural Sciences. 2024;94(3):297-302.
3. Deotale RD, Shinde RD, Meshram SD. Impact of Putrescine and indole-3-butyric acid in enhancing chemical-biochemical parameters and yield of groundnut. Journal of Soils and Crops. 2018;28(2):359-365.
4. Berova, Mand Zlatev Z. Physiological response and yield of paclobutrazol treated tomato plants (*Lycopersicon esculentum* Mill.). Plant Growth Regulation. 2000; 30(2):117-123.
5. Srikanth B, Santhosh B, Rani PS, John K. Effect of Growth Retardants on Plant Yield and Seed Quality of Groundnut (*Arachis hypogaea* L.). Advances in Research. 2024;25(4):57-63.
6. Senoo, Sand Isoda A. Effects of paclobutrazol on dry matter distribution and yield in peanut. Plant production science. 2003;6(1):90-94.
7. Hua S, Zhang Y, Yu H, Lin B, Ding H, Zhang D, Ren Y, Fang Z. Paclobutrazol application effects on plant height seed yield and carbohydrate metabolism in canola. International Journal of Agriculture and Biology. 2014;16:471-479.
8. Barman M, Gunri SK, Puste A, Mand Paul S. Effect of paclobutrazol on growth and yield of kharif groundnut (*Arachis hypogaea* L.). International Journal of Agriculture, Environment and Biotechnology. 2017;10(4):513-518.
9. Gaudin AB. Novel applications in Mississippi cotton and peanut. M.Sc. (Ag.) thesis, State University, Mississippi. 2022; 33(50):5567.
10. Peng Q, Wang H, Tong J, Kabir MH, Huang Z, Xiao L. Effects of indole-3-acetic

acid and auxin transport inhibitor on auxin
distribution and development of peanut at

pegging stage. Scientia horticulturae.
2013;162:76-81.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129027>